

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C11D 3/50, A61K 7/46, C07C 225/14	A1	(11) International Publication Number: WO 00/63339 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/US00/10166 (22) International Filing Date: 14 April 2000 (14.04.00) (30) Priority Data: 60/130,108 20 April 1999 (20.04.99) US (71) Applicant (for all designated States except US): THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): MIRACLE, Gregory, Scot [US/US]; 3065 Old Oxford Road, Hamilton, OH 45013 (US). GRAY, Lon, Montgomery [US/US]; 1806 Promontory Drive, Florence, KY 41042 (US). (74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217-1087 (US).		(81) Designated States: AE, AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), DM, EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PRO-FRAGRANCES		
(57) Abstract		
<p>The present invention relates to fragrance delivery systems which comprise: a) one or more pro-fragrances having formula (I) wherein G¹ and G² are methyl, or a cyclic hydrocarbyl unit derived from the isomers of ionone and/or damascone; R and R¹ are each independently C₁-C₂₂ substituted or unsubstituted, branched or unbranched alkyl, C₂-C₂₂ substituted or unsubstituted, branched or unbranched alkenyl, C₂-C₂₀ substituted or unsubstituted, branched or unbranched hydroxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₃-C₂₀ substituted or unsubstituted cycloalkyl, alkyleneoxy; C₆-C₂₀ aryl, C₅-C₂₀ heteroaryl comprising one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof; two of the units Y¹, Y², R or R¹ can be taken together to form one or more aromatic or non-aromatic, heterocyclic or non-heterocyclic, single rings, fused rings, bicyclo rings, spiroannulated rings, or mixtures thereof, said rings comprising from 3 to 20 carbon atoms and one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof; b) one or more aldehyde releasing oxazolidine pro-fragrances; and c) the balance carriers, pro-fragrances, pro-accords, other perfume ingredients.</p> <div style="text-align: center; margin: 20px 0;"> $\begin{array}{c} \text{R} \quad \text{N} \quad \text{R}^1 \\ \quad \quad \\ \text{G}^1 - \text{CH} - \text{C} - \text{G}^2 \\ \quad \quad \\ \quad \text{R}^2 \quad \text{R}^3 \end{array} \quad (I)$ </div> <div style="text-align: right; margin-top: 50px;"> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

PRO-FRAGRANCES

5

FIELD OF THE INVENTION

The present invention relates to β -amino ketone pro-fragrances which are capable of delivering fragrance raw material ketones, preferably α,β -unsaturated ketones, *inter alia*, damascones, ionones, to a *situs*, especially to human skin wherein said pro-fragrances release the fragrance raw material ketones providing the user with extended aesthetic benefits.

BACKGROUND OF THE INVENTION

Humans have applied scents and fragrances to their skin since antiquity. Originally these aesthetically pleasing materials were commonly isolated in raw form as resins, gums or essential oils from natural sources, *inter alia*, the bark, roots, leaves and fruit of indigenous plants. These resins, gums, and oils were directly applied to the body or diluted with water or other solvent, including in some cases, wine. With the advent of modern chemistry, individual components responsible for the odor properties of these resins, gums and oils were isolated and subsequently characterized. Aside from common "perfume vehicles" *inter alia*, fine perfumes, colognes, eau de toilettes, and after-shave lotions, a wide variety of personal care or personal hygiene items also deliver for aesthetic reasons fragrance notes, accords, or fragrance "characteristics".

It is well known that mixtures of perfume or fragrance raw materials when deposited on the skin lose intensity and may change character with time, mainly due to factors such as differential evaporation and skin penetration. Many attempts have been made to minimize these drawbacks, but so far without notable success. Particularly, efforts have been made to prolong the diffusion, as well as to improve other characteristics of fragrance materials, by e.g. increasing the fragrance raw material concentration or by using additives such as silicones, glycerol, polyethylene glycols and so on. Such additions, however, have never been adequate to increase the longevity of the fragrance odor.

Recently the advent of pro-fragrance and pro-accords have afforded the fine fragrance and perfume formulator with the ability to deliver fragrance raw materials to human skin in a controllable manner thus enhancing the longevity of the fragrance. Most pro-accords *inter alia* orthoesters, are suitable for delivery fragrance raw material alcohols and esters. Aldehydes and ketones have been delivered via acetals and ketals respectively, however, both of these pro-

fragrance materials depend upon the modification of the carbonyl moiety and the rate of ketone fragrance raw material release has been difficult to fine tune to the subtleties of fine fragrance and perfume accords.


Accordingly, there remains a need in the art for a ketone fragrance raw material releasing pro-fragrance which can be formulated into fine fragrances, perfumes, personal care and personal hygiene products wherein the ketone fragrance raw material components can be released in a highly controllable manner to provide enhanced fragrance longevity.

SUMMARY OF THE INVENTION

The present invention meets the aforementioned needs in that it has been surprisingly discovered that certain ketone fragrance raw materials *inter alia* derivatives of ionone and damascone can be controllably released from novel β -amino ketone pro-fragrances. The β -amino pro-fragrances, preferably β -amino ketone pro-fragrances, of the present invention are useful in fragrance delivery systems which deliver fine fragrances with an enhanced perfume longevity.


A first aspect of the present invention relates to fragrance delivery systems comprising:

- A) from about 0.01% by weight, of a pro-fragrance component comprising:
- a) at least 0.01% by weight, of an β -amino ketone pro-fragrance component, said pro-fragrance having the formula:




wherein G^1 and G^2 are methyl, or a unit selected from the group consisting of:


- i) 2,6,6-trimethylcyclohex-2-enyl having the formula:




- ii) 2,6,6-trimethylcyclohex-1-enyl having the formula:



iii) 2-methylene-6,6-dimethylcyclohexanyl having the formula:




iv) 2,6,6-trimethylcyclohex-3-enyl having the formula:




v) and mixtures thereof;

provided at least one G^1 or G^2 is methyl and further provided both G^1 and G^2 are not both methyl; R and R^1 are each independently C_1 - C_{22} substituted or unsubstituted, branched or unbranched alkyl, C_2 - C_{22} substituted or unsubstituted, branched or unbranched alkenyl, C_2 - C_{20} substituted or unsubstituted, branched or unbranched hydroxyalkyl, C_7 - C_{20} substituted or unsubstituted alkylenearyl, C_3 - C_{20} substituted or unsubstituted cycloalkyl, alkyleneoxy units having the formula:



wherein R^4 is C_2 - C_4 alkylene, R^5 is hydrogen, C_1 - C_4 alkyl, and mixtures thereof, x is from 1 to 6; C_6 - C_{20} aryl, C_5 - C_{20} heteroaryl comprising one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof; two of the units Y^1 , Y^2 , R or R^1 can be taken together to form one or more aromatic or non-aromatic, heterocyclic or non-heterocyclic, single rings, fused rings, bicyclo rings, spiroannulated rings, or mixtures thereof, said rings comprising from 3 to 20 carbon atoms and one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof;

- b) at least 0.01% by weight, of an aldehyde or ketone releasing pro-fragrance component, said pro-fragrance having the formula:



wherein said pro-fragrance or pro-accord releases an aldehyde or a ketone fragrance raw material, wherein X is oxygen or sulfur; R is:


- i) C₆-C₂₂ substituted or unsubstituted linear alkyl;
- ii) C₆-C₂₂ substituted or unsubstituted branched alkyl;
- iii) C₆-C₂₂ substituted or unsubstituted linear alkenyl;
- iv) C₆-C₂₂ substituted or unsubstituted branched alkenyl;
- v) C₆-C₂₂ substituted or unsubstituted cycloalkyl;
- vi) C₆-C₂₂ substituted or unsubstituted branched cycloalkyl;
- vii) C₆-C₂₂ substituted or unsubstituted cycloalkenyl;
- viii) C₆-C₂₂ substituted or unsubstituted branched cycloalkenyl;
- ix) C₆-C₂₂ substituted or unsubstituted aryl;
- x) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
- xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
- xii) and mixtures thereof;

R¹ is:

- i) hydrogen;
- ii) C₁-C₁₀ substituted or unsubstituted linear alkyl;
- iii) C₃-C₁₀ substituted or unsubstituted branched alkyl;
- iv) C₂-C₁₀ substituted or unsubstituted linear alkenyl;
- v) C₃-C₁₀ substituted or unsubstituted branched alkenyl;
- vi) C₃-C₁₅ substituted or unsubstituted cycloalkyl;
- vii) C₄-C₁₅ substituted or unsubstituted branched cycloalkyl;
- viii) C₄-C₁₅ substituted or unsubstituted cycloalkenyl;
- ix) C₅-C₁₅ substituted or unsubstituted branched cycloalkenyl;
- x) C₆-C₁₅ substituted or unsubstituted aryl;
- xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
- xii) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;

R and R¹ can be taken together to form a substituted or unsubstituted ring having in the ring from 3 to 10 carbon atoms; and each R², R³, R⁶ and each R⁴ and R⁵ pair are independently:

- i) R¹;
- ii) hydroxyl;
- iii) a carbonyl comprising unit having the formula:



wherein R⁷ is:

- a) -OH;
 - b) -OR⁸ wherein R⁸ is hydrogen, C₁-C₁₅ substituted or unsubstituted linear alkyl, C₁-C₁₅ substituted or unsubstituted branched alkyl, C₂-C₂₂ substituted or unsubstituted linear alkenyl, C₃-C₂₂ substituted or unsubstituted branched alkenyl, or mixtures thereof;
 - c) -N(R⁹)₂ wherein R⁹ is hydrogen, C₁-C₆ substituted or unsubstituted linear alkyl, C₃-C₆ substituted or unsubstituted branched alkyl, or mixtures thereof;
 - d) C₁-C₂₂ substituted or unsubstituted linear alkyl;
 - e) C₁-C₂₂ substituted or unsubstituted branched alkyl;
 - f) C₂-C₂₂ substituted or unsubstituted linear alkenyl;
 - g) C₃-C₂₂ substituted or unsubstituted branched alkenyl;
 - h) C₃-C₂₂ substituted or unsubstituted cycloalkyl;
 - i) C₆-C₂₂ substituted or unsubstituted aryl;
 - j) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 - k) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
- the index x is from 0 to 22;
- iv) alkyleneoxy units having the formula:



wherein each R¹⁰, R¹¹, and R¹² is independently;

- a) hydrogen;
- b) -OH;
- c) C₁-C₄ alkyl;
- d) or mixtures thereof;

R¹³ is:

- a) hydrogen;
- b) C₁-C₄ alkyl;
- c) or mixtures thereof;

R¹⁴ is:

- a) hydrogen;
- b) C₁-C₄ alkyl;
- c) or mixtures thereof;

R¹⁰ and R¹¹ can be taken together to form a C₃-C₆ spiroannulated ring, carbonyl unit, or mixtures thereof; y has the value from 0 to 10, z has the value from 1 to 50;

v) any two R², R³, R⁴, R⁵, or R⁶ units can be taken together to form:

- a) a carbonyl moiety;
- b) a C₃-C₆ spiroannulated ring;
- c) a heterocyclic aromatic ring comprising from 5 to 7 atoms;
- d) a non-heterocyclic aromatic ring comprising from 5 to 7 atoms;
- e) a heterocyclic ring comprising from 5 to 7 atoms;
- f) a non-heterocyclic ring comprising from 5 to 7 atoms;
- g) or mixtures thereof;

vi) and mixtures thereof; and

the index n is an integer from 1 to 3; and

B) optionally from about 1% by weight, a fragrance raw material component comprising:

- i) optionally at least 1% by weight, of a mixture of one or more base note fragrances;
- ii) optionally at least 1% by weight, of a mixture of one or more top or middle note fragrances;
- iii) optionally the balance carriers, fixatives, and other adjunct ingredients.


These and other objects, features and advantages will become apparent to those of ordinary skill in the art from a reading of the following detailed description and the appended claims. All percentages, ratios and proportions herein are by weight, unless otherwise specified. All temperatures are in degrees Celsius (°C) unless otherwise specified. All documents cited are

in relevant part, incorporated herein by reference.

DETAILED DESCRIPTION OF THE INVENTION


The present invention relates to fragrance delivery systems which comprise one or more β -amino ketone pro-fragrances of damascones, ionones and their isomers. The pro-fragrances of the present invention are suitable for use in fragrance delivery systems which are capable in delivering enhanced fragrance benefits to fine fragrances and perfumes as well as personal care and personal hygiene articles.

The β -amino pro-fragrances of the present invention have the formula:




wherein G^1 and G^2 are methyl, or a unit selected from the group consisting of:


- i) 2,6,6-trimethylcyclohex-2-enyl having the formula:




- ii) 2,6,6-trimethylcyclohex-1-enyl having the formula:



- iii) 2-methylene-6,6-dimethylcyclohexanyl having the formula:



- iv) 2,6,6-trimethylcyclohex-3-enyl having the formula:



v) and mixtures thereof;

provided at least one G^1 or G^2 is methyl and further provided both G^1 and G^2 are not both methyl.

R and R^1 are each independently C_1 - C_{22} substituted or unsubstituted, branched or
 5 unbranched alkyl, C_2 - C_{22} substituted or unsubstituted, branched or unbranched alkenyl, C_2 - C_{20}
 substituted or unsubstituted, branched or unbranched hydroxyalkyl, C_7 - C_{20} substituted or
 unsubstituted alkylenearyl, C_3 - C_{20} substituted or unsubstituted cycloalkyl, , alkyleneoxy units
 having the formula:




10 wherein R^4 is C_2 - C_4 alkylene, R^5 is hydrogen, C_1 - C_4 alkyl, and mixtures thereof, x is from 1 to 6;
 C_6 - C_{20} aryl, C_5 - C_{20} heteroaryl comprising one or more heteroatoms selected from the group
 consisting of nitrogen, oxygen, sulfur, and mixtures thereof; two of the units Y^1 , Y^2 , R or R^1 can
 be taken together to form one or more aromatic or non-aromatic, heterocyclic or non-
 heterocyclic, single rings, fused rings, bicyclo rings, spiroannulated rings, or mixtures thereof,
 15 said rings comprising from 3 to 20 carbon atoms and one or more heteroatoms selected from the
 group consisting of nitrogen, oxygen, sulfur, and mixtures thereof.

The following are the common chemical names for the damascones and ionones which
 are delivered by the pro-fragrances of the present invention; 1-(2,6,6-trimethylcyclohex-2-enyl)-
 2-butene-1-one (α -damascone), 1-(2,6,6-trimethylcyclohex-1-enyl)-2-butene-1-one (β -
 20 damascone), 1-(2-methylene-6,6-dimethylcyclohexanyl)-2-butene-1-one (γ -damascone), 1-
 (2,6,6-trimethylcyclohex-3-enyl)-2-butene-1-one (δ -damascone), 4-(2,6,6-trimethylcyclohex-2-
 enyl)-3-butene-2-one (α -ionone), 4-(2,6,6-trimethylcyclohex-1-enyl)-3-butene-2-one (β -ionone),
 4-(2-methylene-6,6-dimethylcyclohexanyl)-3-butene-2-one (γ -ionone).


Further preferred ketones which can be delivered by the β -amino ketone pro-fragrances
 25 of the present invention include 4-(2,6,6-trimethyl-2-cyclohexen-1-yl)-3-methyl-3-buten-2-one,
 1-(2,6,6-trimethyl-2-cyclohexen-1-yl)-1,6-heptadien-3-one, 4-(2,5,6,6-tetramethyl-2-
 cyclohexen-1-yl)-3-methyl-3-buten-2-one, 2-pentyl-2-cyclopenten-1-one, 2-hexyl-2-cyclopenten-
 1-one, *cis*-3-methyl-2-(2-pentenyl)-2-cyclopenten-1-one, 6-(2-propenyl)-5-methyl-2-(1-
 methylethylidene)cyclohexanone, 4-(4-methoxy)phenyl-3-buten-2-one, 2,5,5-trimethyl-2,6-
 30 heptadien-4-one, and 2-methyl-6-(4-methylcyclohex-3-enyl)-2,5-heptadien-4-one.

An example of a β -amino ketone pro-fragrance according to the present invention includes 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N-(2-hydroxyethyl)-N-phenylmethyl-1-butanone having the formula:




5

1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-di(2-hydroxyethyl)-1-butanone having the formula:




1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone having the formula:



10

and 2-hexyl-3-di(2-hydroxyethyl)aminocyclopentanone having the formula:




Aldehyde and Ketone Releasing Pro-fragrances


The fragrance delivery systems of the present invention comprise one or more heterocyclic aldehyde-releasing and/or ketone-releasing pro-fragrances, preferably oxazolidines, tetrahydro-1,3-oxazines, thiazolidines, or tetrahydro-1,3-thiazines, more preferably oxazolidines, or tetrahydro-1,3-oxazines, most preferably oxazolidines.

15


The pro-fragrances or pro-accords which are suitable for use in the fragrance delivery systems described herein have the formula:



- 5 and are capable of releasing an aldehyde fragrance raw material having the formula:



or a ketone fragrance raw material having the formula:



10

For the purposes of the fragrance delivery systems which comprise one or more heterocyclic pro-fragrances or pro-accords, R units are defined herein as:

- i) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted linear alkyl; one or more examples of a fragrance raw material which comprises this unit includes nonanal and decanal;
- 15 ii) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted branched alkyl; one or more examples of a fragrance raw material which comprises this unit includes 2-methyldecanal;
- 20 iii) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted linear alkenyl; one or more examples of a fragrance raw material which comprises this unit includes 10-undecenal;
- iv) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted branched alkenyl; one or more examples of a fragrance raw material which comprises this unit includes citral, melonal, and neral;
- 25 v) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₀ substituted or unsubstituted cycloalkyl; one or more examples of a fragrance raw material which comprises this unit includes cyclopentadecanone;

- vi) C₆-C₂₂, preferably C₆-C₁₅, more preferably C₆-C₁₅ substituted or unsubstituted branched cycloalkyl; one or more examples of a fragrance raw material which comprises this unit includes camphor and muscone;
- 5 vii) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted cycloalkenyl; one or more examples of a fragrance raw material which comprises this unit includes civetone;
- viii) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted branched cycloalkenyl; one or more examples of a fragrance raw material which comprises this unit includes α-damascone and β-ionone;
- 10 ix) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₀ substituted or unsubstituted aryl wherein said aryl unit preferably comprises a phenyl unit; one or more examples of a fragrance raw material which comprises this unit includes benzaldehyde, hydrotropaldehyde and vanillin;
- x) C₆-C₂₂, preferably C₆-C₁₅, more preferably C₆-C₁₅ substituted or unsubstituted heterocyclicalkyl;
- 15 xi) C₆-C₂₂, preferably C₆-C₁₈, more preferably C₆-C₁₅ substituted or unsubstituted heterocyclicalkenyl; one or more examples of a fragrance raw material which comprises this unit includes;
- xii) and mixtures thereof;
- 20 R¹ is:
- i) hydrogen;
- ii) C₁-C₁₀, preferably C₁-C₅ substituted or unsubstituted linear alkyl;
- iii) C₃-C₁₀, preferably C₃-C₅ substituted or unsubstituted branched alkyl;
- iv) C₂-C₁₀, preferably C₂-C₅ substituted or unsubstituted linear alkenyl;
- 25 v) C₃-C₁₀, preferably C₄-C₁₀ substituted or unsubstituted branched alkenyl;
- vi) C₃-C₁₅, preferably C₆-C₁₀ substituted or unsubstituted cycloalkyl;
- vii) C₄-C₁₅, preferably C₆-C₁₀ substituted or unsubstituted branched cycloalkyl;
- viii) C₄-C₁₅, preferably C₆-C₁₀ substituted or unsubstituted cycloalkenyl;
- ix) C₅-C₁₅, preferably C₆-C₁₀ substituted or unsubstituted branched cycloalkenyl;
- 30 x) C₆-C₁₅, preferably C₆-C₁₀ substituted or unsubstituted aryl;
- xi) C₆-C₂₂, preferably C₆-C₁₀ substituted or unsubstituted heterocyclicalkyl;
- xii) C₆-C₂₂, preferably C₆-C₁₀ substituted or unsubstituted heterocyclicalkenyl;
- R and R¹ can be taken together to form a substituted or unsubstituted ring having in the ring from 3 to 10 carbon atoms; one or more examples of which are substituted
- 35 cyclopentanone derivatives inter alia hedione and nectaryl; and

each R^2 , R^3 , R^6 and each R^4 and R^5 pair are independently:

- i) R^1 ;
- ii) hydroxyl;
- iii) a carbonyl comprising unit having the formula:



wherein R^7 is:

- a) -OH, in the case of carboxylic acids;
- b) -OR⁸, in the case of esters wherein R^8 is hydrogen; C_1 - C_{15} , preferably C_1 - C_{10} , more preferably C_1 - C_4 substituted or unsubstituted linear alkyl; C_3 -
10 C_{15} , preferably C_3 - C_{10} , more preferably C_3 - C_4 substituted or unsubstituted branched alkyl; C_2 - C_{22} , preferably C_2 - C_{10} , more preferably C_2 - C_4 substituted or unsubstituted linear alkenyl; C_3 - C_{22} substituted or unsubstituted branched alkenyl, or mixtures thereof;
- c) -N(R^9)₂ in the case of amides wherein each R^9 is independently
15 hydrogen; C_1 - C_{15} , preferably C_1 - C_{10} , more preferably C_1 - C_4 substituted or unsubstituted linear alkyl; C_3 - C_{15} , preferably C_3 - C_{10} , more preferably C_3 - C_4 substituted or unsubstituted branched alkyl; or mixtures thereof;
- d) C_1 - C_{22} , preferably C_1 - C_5 substituted or unsubstituted linear alkyl;
- e) C_1 - C_{22} , preferably C_3 - C_5 substituted or unsubstituted branched alkyl;
- 20 f) C_2 - C_{22} , preferably C_2 - C_5 substituted or unsubstituted linear alkenyl;
- g) C_3 - C_{22} , preferably C_4 - C_{10} substituted or unsubstituted branched alkenyl;
- h) C_5 - C_{22} , preferably C_6 - C_{10} substituted or unsubstituted cycloalkyl;
- i) C_6 - C_{22} , preferably C_6 - C_{10} substituted or unsubstituted aryl;
- j) C_6 - C_{22} , preferably C_6 - C_{10} substituted or unsubstituted heterocyclicalkyl;
- 25 k) C_6 - C_{22} , preferably C_6 - C_{10} substituted or unsubstituted heterocyclicalkenyl;

the index is from 0 to 22;

- v) alkyleneoxy units having the formula:



wherein each R^{10} , R^{11} , and R^{12} is independently;

- a) hydrogen;
- b) -OH;
- c) C_1 - C_4 alkyl, preferably methyl;

d) or mixtures thereof; preferably R^{10} , R^{11} , and R^{12} are each hydrogen;

R^{13} is:

a) hydrogen;

b) C_1 - C_4 alkyl, preferably methyl;

c) or mixtures thereof; preferably R^{13} is methyl or hydrogen, more preferably hydrogen;

R^{14} is:

a) hydrogen;

b) C_1 - C_4 alkyl, preferably methyl;

c) or mixtures thereof; preferred R^{14} is hydrogen;

R^{10} and R^{11} can be taken together to form a C_3 - C_6 spiroannulated ring, carbonyl unit, or mixtures thereof; y has the value from 0 to 10, z has the value from 1 to 50;

vi) any two R^2 , R^3 , R^4 , R^5 , or R^6 units can be taken together, and where feasible, combined to form:

a) a carbonyl moiety;

b) a C_3 - C_6 spiroannulated ring;

c) a heterocyclic aromatic ring comprising from 5 to 7 atoms;

d) a non-heterocyclic aromatic ring comprising from 5 to 7 atoms;

e) a heterocyclic ring comprising from 5 to 7 atoms;

f) a non-heterocyclic ring comprising from 5 to 7 atoms;

g) or mixtures thereof;

vii) and mixtures thereof;

the index n is an integer from 1 to 3, preferably 1 or 2, more preferably 1.

For the purposes of the present invention, the term "substituted" is defined herein as "compatible moieties which replace a hydrogen atom". For the purposes of the present invention, hydrogens which are substitutable are labeled as R' units in the following examples. Non-limiting examples of substituents which can replace hydrogen atoms are C_1 - C_{22} linear or branched hydrocarbyl units *inter alia* alkyl, alkenyl; hydroxy, nitrilo, nitro, carboxyl (-CHO; -CO₂H; -CO₂R"; -CONH₂; -CONHR"; -CONR"2; wherein R" is C_1 - C_{12} linear or branched alkyl), amino, C_1 - C_{12} mono- and dialkylamino, and mixtures thereof. However, the formulator may wish to include other substituents not specifically mentioned herein. Not each hydrogen of a substituted unit, i.e., substituted linear alkyl, must be substituted; only one hydrogen must be substituted by another moiety for a unit to be "substituted" for the purposes of the present invention.

FRAGRANCE DELIVERY SYSTEMS

The present invention also relates to fragrance raw material delivery systems, said delivery systems comprising:

- 5 A) from about 0.1%, preferably from about 0.5%, more preferably from about 1%, yet more preferably from about 5%, most preferably from about 10% to about 100%, preferably to about 75%, more preferably to about 50% by weight, of a pro-fragrance component comprising:
- 10 i) at least 0.01%, preferably from about 0.5%, more preferably from about 5%, most preferably from about 25% to about 100%, more preferably to about 75%, most preferably to about 50% by weight, of an β -amino ketone pro-fragrance component according to the present invention;
- 15 ii) optionally from about 0.01%, preferably from about 0.5%, more preferably from about 5%, most preferably from about 25% to about 99%, more preferably to about 75%, most preferably to about 50% by weight, by weight, of one or more pro-accords formed from at least one fragrance raw material, wherein said pro-accord is selected from the group consisting of acetals, ketals, orthoesters, orthocarbonates, and mixtures thereof, each pro-accord releasing upon hydrolysis said
- 20 fragrance raw material from which it is formed, said fragrance raw materials selected from the group consisting of primary, secondary, and tertiary alcohols, aldehydes, ketones, esters, carbonates, and mixtures thereof, provided at least one pro-accord:
- 25 a) is formed from at least one fragrance raw material having a molecular weight greater than or equal to about 100 g/mol;
- b) has a fragrance release half-life of greater than or equal to about 0.1 hours at pH 5.3 and less than or equal to about 12 hours at pH 2.5 when measured in NaH_2PO_4 buffer;
- 30 iii) the balance carriers, stabilizers, and other adjunct ingredients; and
- B) optionally from about 0.01%, preferably from about 0.5%, more preferably from about 1%, yet more preferably from about 5%, most preferably from about 10% to about 99%, preferably to about 75%, more preferably to about 50% by weight, by weight, a fragrance raw material component comprising:
- 35 i) optionally at least 0.01%, preferably from about 0.5%, more preferably from about 5%, most preferably from about 25% to about 100%, more

preferably to about 75%, most preferably to about 50% by weight, by weight, of a mixture of one or more base note fragrances;

- ii) optionally at least 0.01%, preferably from about 0.5%, more preferably from about 5%, most preferably from about 25% to about 100%, more preferably to about 75%, most preferably to about 50% by weight, by weight, of a mixture of one or more top or middle note fragrances;

- iii) optionally the balance carriers, fixatives, and other adjunct ingredients.

When used the fragrance delivery systems of the present invention will comprise from about 0.1%, preferably from about 1% to about 100%, preferably to about 85% of the composition depending upon the execution. For example, a fine fragrance may comprise 100% by weight, of a fragrance delivery as described herein, whereas a body lotion or cream may comprise from about 1% to about 5% by weight, of the fragrance delivery system.

For the purposes of the fragrance delivery systems of the present invention, a "pro-accord which comprises n fragrance raw materials but which releases $n+1$ fragrance raw materials" is defined as "a compound which is prepared from one or more fragrance raw materials, said fragrance raw material being chemically transformed into a "releasable form" such that when said releasable form breaks down, the original fragrance raw material is released as well as at least one other fragrance raw material which was not a starting material used in forming the releasable form". The term "releasable form" is defined herein as a "pro-fragrance or pro-accord compound, which ever form is applicable". Non-limiting examples of "releasable forms" or pro-accords which satisfy the $n + 1$ requirement are as follows.

Preferably the pro-accords of the present invention are orthoesters and orthocarbonates which deliver $n + 1$ fragrance raw materials when the "pro-accord" has been formed from n fragrance raw materials. These "Increased Release" pro-accords are preferably formate, acetate, propionate, and benzoate orthoesters. Any fragrance raw material may be used to form the "increased release" pro-accord provided the final pro-accord:


- a) has a molecular weight greater than or equal to about 300 g/mol;
- b) has a molecular weight at least two times greater than the lowest molecular weight fragrance raw material which comprises said pro-accord; and
- c) has a fragrance release half-life greater than or equal to about 0.1 hours when measured in NaH_2PO_4 buffer at pH 5.3 and less than about 12 hours when measured in NaH_2PO_4 buffer at pH 2.5.

The value of the index n is an integer from 1 to 3.


When present the "increased release" pro-accords comprise at least 0.01% of the increased fragrance retention composition, preferably at least about 0.1% more preferably at least

about 0.5% by weight, of said composition. More than one "increased release" pro-accords may be combined together as described herein above to provide a complex perfume mixture or accord.

- Non-limiting examples of pro-accords which optionally comprise the fragrance delivery systems of the present invention include orthoesters having the formula:



- wherein R is hydrogen, C₁-C₈ linear alkyl, C₄-C₂₀ branched alkyl, C₆-C₂₀ cyclic alkyl, C₆-C₂₀ branched cyclic alkyl, C₆-C₂₀ linear alkenyl, C₆-C₂₀ branched alkenyl, C₆-C₂₀ cyclic alkenyl, C₆-C₂₀ branched cyclic alkenyl, C₆-C₂₀ substituted or unsubstituted aryl, and mixtures thereof;
- R¹, R² and R³ are independently C₁-C₂₀ linear, branched, or substituted alkyl; C₂-C₂₀ linear, branched, or substituted alkenyl; C₅-C₂₀ substituted or unsubstituted cyclic alkyl; C₆-C₂₀ substituted or unsubstituted aryl, C₂-C₄₀ substituted or unsubstituted alkyleneoxy; C₃-C₄₀ substituted or unsubstituted alkyleneoxyalkyl; C₆-C₄₀ substituted or unsubstituted alkylenearyl; C₆-C₃₂ substituted or unsubstituted aryloxy; C₆-C₄₀ substituted or unsubstituted alkyleneoxyaryl; C₆-C₄₀ oxyalkylenearyl, and mixtures thereof; or a cyclic orthoester having the formula:



- wherein at least one R⁴ or R⁵ is derived from a fragrance raw material alcohol, each X is -C(R⁶)₂- wherein each R⁶ is independently hydrogen, C₁-C₂₂ linear or branched alkyl, C₂-C₂₂ linear or branched alkenyl, C₆-C₂₂ substituted or unsubstituted aryl, and mixtures thereof, Y is -CR⁷R⁸-, C=O, and mixtures thereof, wherein R⁷ and R⁸ are independently hydrogen, hydroxyl, nitro, nitrilo, C₁-C₃₀ substituted or unsubstituted linear alkyl, C₃-C₃₀ substituted or unsubstituted branched alkyl, C₃-C₃₀ substituted or unsubstituted cyclic alkyl, C₂-C₃₀ substituted or unsubstituted linear alkenyl, C₃-C₃₀ substituted or unsubstituted branched alkenyl, C₃-C₃₀ substituted or unsubstituted cyclic alkenyl, C₂-C₃₀ substituted or unsubstituted linear alkynyl, C₃-C₃₀ substituted or unsubstituted branched alkynyl, C₆-C₃₀ substituted or unsubstituted alkylenearyl, C₆-C₃₀ substituted or unsubstituted aryl, C₂-C₂₀ substituted or unsubstituted alkyleneoxy, C₃-C₂₀ substituted or unsubstituted alkyleneoxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₆-C₂₀ substituted or unsubstituted alkyleneoxyaryl, and mixtures

thereof, or R⁷ and R⁸ can be taken together to form a spiroannulated ring or taken together with any R⁶ to form a fused ring, said spiroannulated or fused ring having from 3 to 8 carbons and optionally one or more heteroatoms in said ring, said ring further optionally substituted by one or more C₁-C₂₂ alkyl, C₁-C₂₂ alkenyl, C₆-C₁₂ aryl, C₆-C₂₂ alkylenearyl units, and mixtures thereof; m is from 0 to 14, p is from 0 to 14, and n is from 0 to 3; provided m + n + p is at least 1 and less than or equal to 14.

Examples of "increased release" pro-accords which are comprised of a single fragrance raw materials but which release at least 2 fragrance raw materials include but are not limited to tris(citronellyl) orthoformate which releases the binary accord citronellyl formate/citronellol; tris(citronellyl) orthoacetate which releases the binary accord citronellol/citronellyl acetate; tris(geranyl) orthoformate which releases the binary accord geraniol/geranyl formate; tris(geranyl) orthoacetate which releases the binary accord geraniol/geranyl acetate; tris(phenylethyl) orthoformate which releases the binary accord 2-phenylethanol/phenylethyl formate; tris(phenylethyl) orthoacetate which releases the binary accord 2-phenylethanol/phenylethyl acetate; tris(9-decenyl) orthoformate which releases the binary accord 9-decen-1-ol/9-decenyl formate; and tris(9-decenyl) orthoacetate which releases the binary accord 9-decen-1-ol/9-decenyl acetate. For the purposes of the present invention when an orthoester comprises only one fragrance raw material alcohol (releases a binary accord) the terms "tris(alcohol) orthoester" and "alcohol orthoester" are used interchangeably. For example, tris(geranyl) orthoacetate stands equally well for geranyl orthoacetate and are considered interchangeable.

The pro-accord tris(9-decenyl) orthoformate is prepared by treating 9-decenol (i.e., Rosalva), which is a fragrance raw material as defined herein, with a suitable amount of triethyl orthoformate, not a fragrance raw material as defined herein, in the presence of an acid catalyst optionally in the presence of a solvent. Tris(9-decenyl) when exposed to suitable conditions (e.g., exposure to the acid mantle of human skin) breaks down to release a mixture of 9-decenol and 9-decenyl formate, both of which are fragrance raw materials. Therefore, one fragrance raw material is used to prepare a releasable form (pro-accord) of two fragrance raw materials.

The pro-accord 3,7-dimethyl-1,6-octadien-3-yl 3-(β-naphthyl)-3-oxo-propionate, which is a β-ketoester pro-accord, is prepared by treating 3,7-dimethyl-1,6-octadien-3-ol (linalool), which is a fragrance raw material according to the present invention, with diketene under suitable conditions to form intermediate 3,7-dimethyl-1,6-octadien-3-yl 3-oxo-butyrate, which is subsequently treated with 2-naphthoyl chloride to yield the pro-accord. 3,7-Dimethyl-1,6-octadien-3-yl 3-(β-naphthyl)-3-oxo-propionate when exposed to suitable conditions (e.g.,

exposure to nascent moisture) breaks down to release a mixture of linalool and methyl β -naphthyl ketone, both of which are fragrance raw materials as defined herein.

As described herein above, an optional component of the fragrance delivery systems of the present invention are pro-fragrances or pro-accords which are not heterocyclic aldehyde and/or ketone releasing pro-fragrances. The optional pro-accords or pro-fragrances are equally functional in either personal care compositions *inter alia* lotions, creams, deodorants or personal fragrance compositions *inter alia* fine fragrances, perfumes.

Preferred optional pro-accords and/or pro-fragrances include, but are not limited to, orthoesters, orthocarbonates, acetals, ketals, ortholactones, and β -ketoesters.

Non-limiting examples of optional orthoesters which are suitable for use in the fragrance delivery systems of the present invention include tris-geranyl orthoformate, tris(*cis*-3-hexen-1-yl) orthoformate, tris(phenylethyl) orthoformate, bis(citronellyl) ethyl orthoacetate, tris(citronellyl) orthoformate, tris(*cis*-6-nonenyl) orthoformate, tris(phenoxyethyl) orthoformate, tris(geranyl, neryl) orthoformate (70:30 geranyl:neryl), tris(9-decenyl) orthoformate, tris(3-methyl-5-phenylpentanyl) orthoformate, tris(6-methylheptan-2-yl) orthoformate, tris([4-(2,2,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-yl] orthoformate, tris[3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-yl] orthoformate, trismenthyl orthoformate, tris(4-isopropylcyclohexylethyl-2-yl) orthoformate, tris-(6,8-dimethylnonan-2-yl) orthoformate, tris-phenylethyl orthoacetate, tris(*cis*-3-hexen-1-yl) orthoacetate, tris(*cis*-6-nonenyl) orthoacetate, tris-citronellyl orthoacetate, bis(geranyl) benzyl orthoacetate, tris(geranyl) orthoacetate, tris(4-isopropylcyclohexylmethyl) orthoacetate, tris(benzyl) orthoacetate, tris(2,6-dimethyl-5-heptenyl) orthoacetate, bis(*cis*-3-hexen-1-yl) amyl orthoacetate, and neryl citronellyl ethyl orthobutyrate.

Non-limiting examples of optional orthocarbonates which are suitable for use in the fragrance delivery systems of the present invention include bis(ethyl) bis(geranyl) orthocarbonate, bis(ethyl) bis(phenylethyl) orthocarbonate, bis(ethyl) bis(*cis*-3-hexenyl) orthocarbonate, bis(ethyl) bis(citronellyl) orthocarbonate, bis(ethyl) bis(linalyl) orthocarbonate, bis(ethyl) bis(menthyl) orthocarbonate, bis(dodecyl) bis(geranyl) orthocarbonate, and bis(dodecyl) bis(phenylethyl) orthocarbonate.

Non-limiting examples of optional acetals which are suitable for use in the fragrance delivery systems of the present invention include bis(*cis*-3-hexenyl) vanillin, bis(geranyl) cinnamaldehyde acetal, bis(2-phenylethyl) anisaldehyde acetal, bis(citronellyl) cyclamen aldehyde acetal, and bis(citronellyl) citral acetal.

Non-limiting examples of optional ketals which are suitable for use in the fragrance delivery systems of the present invention include bis(linalyl) β -ionone ketal,

bis(dihydromyrcenyl) α -damascone ketal, bis(linalyl) 6,7-dihydro-1,1,2,3,3-pentamethyl-4(5H)-indanone ketal, bis(dihydromyrcenyl) β -ionone ketal, and bis(citronellyl) *cis*-jasmane ketal.

Non-limiting examples of optional β -ketoesters which are suitable for use in the fragrance delivery systems of the present invention include 2,6-dimethyl-7-octen-2-yl 3-(4-methoxyphenyl)-3-oxo-propionate, 3,7-dimethyl-1,6-octadien-3-yl 3-(α -naphthyl)-3-oxo-propionate, 2,6-dimethyl-7-octen-2-yl 3-(4-methoxyphenyl)-3-oxo-propionate, *cis* 3-hexen-1-yl 3-(β -naphthyl)-3-oxo-propionate, 2,6-dimethyl-7-octen-2-yl 3-(nonanyl)-3-oxo-propionate, 2,6-dimethyl-7-octen-2-yl 3-oxo-butyrate, 3,7-dimethyl-1,6-octadien-3-yl 3-oxo-butyrate, 2,6-dimethyl-7-octen-2-yl 3-(β -naphthyl)-3-oxo-2-methylpropionate, 3,7-dimethyl-1,6-octadien-3-yl 3-(β -naphthyl)-3-oxo-2,2-dimethylpropionate, 3,7-dimethyl-1,6-octadien-3-yl 3-(β -naphthyl)-3-oxo-2-methylpropionate, 3,7-dimethyl-2,6-octadienyl 3-(β -naphthyl)-3-oxo-propionate, and 3,7-dimethyl-2,6-octadienyl 3-heptyl-3-oxo-propionate.

An example of a fragrance delivery system according to the present invention comprises:

- a) from about 0.1% by weight, of a β -amino ketone pro-fragrance, for example, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone;
- b) from about 0.2% by weight, of one or more aldehyde releasing pro-fragrances, for example, 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-oxazikudubecarboxylic acid methyl ester, 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester;
- c) optionally from about 0.1% by weight, of one or more alcohol releasing pro-fragrances; preferably an alcohol releasing pro-accord, for example, geranyl orthoformate;
- d) from about 0.1% by weight, of one or more fragrance raw materials, for example, lillial, γ -decalactone;
- e) optionally, from about 1% by weight, of pre-blended perfume ingredients of fragrance raw material accords; and
- f) the balance carriers.

A non-limiting example of an alcohol releasing pro-accord admixture which is suitable for use in the fragrance delivery systems of the present invention comprises:


- i) from about 0.5% to about 5% by weight, of nonadyl orthoformate;
- ii) from about 1.5% to about 5% by weight, of mugetanol orthoformate;
- iii) from about 1.5% to about 5% by weight, of osyrol orthoformate; and
- iv) from about 1% to about 10% by weight, of benzyl orthoacetate.

Fragrance Release Half-life

One aspect of the present invention which is a key element in providing the formulator with a method for determining the manner in which a pro-fragrance according to the present invention releases its fragrance raw material, is the measurement of the pro-fragrance "Fragrance Release Half-Life, (FRHL). The pro-fragrances useful in the personal care compositions of the present invention generally have a delayed release of final fragrance raw material in order to achieve the increased fragrance longevity benefits described herein. However, the pro-fragrances generally also deliver the fragrance raw materials during a time period useful to the formulator, for example, within a time period desirable to the consumer.

For the purposes of the present invention the pro-accords generally have a FRHL of less than or equal to 12 hours when measured in NaH_2PO_4 buffer at pH 2.5 and greater than or equal to 0.1 hour when measured in NaH_2PO_4 buffer at pH 5.3. The "Fragrance Release Half-life" is defined herein as follows.

Pro-fragrances deliver their corresponding mixture of fragrance raw materials or fragrance accords according to the equation:



wherein the fragrance raw material which is released may be released as a single component or a multiple fragrance raw material accord.

The rate at which the fragrance is released is defined by the formula:

$$\text{Rate} = k[\text{Pro-fragrance}]$$

and can be further expressed by the formula:

$$-\frac{d[\text{Pro-fragrance}]}{dt} = k[\text{Pro-fragrance}]$$

wherein k is the release rate constant and $[\text{Pro-fragrance}]$ is the concentration of pro-fragrance.

For the purposes of the present invention the "Fragrance Release Half-life", $t_{1/2}$, is related to the release rate constant by the formula:

$$t_{1/2} = \frac{0.693}{k}$$

and this relationship is used for the purposes of the present invention to determine the "FRHL.

Due to the hydrophobic nature of some pro-accords, it is necessary to conduct the determination of $t_{1/2}$ and k in a mixture of 90/10 dioxane/phosphate buffered water. The

phosphate buffered water is prepared by admixing 3.95 mL of 85% phosphoric acid (H_3PO_4) and 24 g of sodium dihydrogen phosphate (NaH_2PO_4) with one liter of water. The pH of this solution is approximately 2.5. Next 10 mL of the phosphate buffer is admixed with 90 mL of dioxane and the pro-fragrance to be analyzed is added. The hydrolysis kinetics are then
5 monitored by conventional HPLC at 30° C.

In some instances, it is desirable to formulate a fragrance delivery system having one or more pro-fragrances which deliver a rapid release of fragrance raw material in addition to the delayed onset of a fragrance. In such cases the hydrolysis rate, and therefore the determination of $t_{1/2}$ must be measured in a buffer system which can accommodate this more rapid hydrolysis
10 rate.

The pro-fragrances of the present invention are stable under pH conditions encountered in the formulation and storage of fine perfume, personal care and personal hygiene articles which have a pH of from about 7.1 to 11.5, and during solution-use of such products. Due to their high molecular weight and hydrophobicity, these pro-fragrances and/or pro-accords remain deposited
15 upon skin even when exposed to water (i.e. when formulated into a sun screen). Because the pro-fragrances are subject to hydrolysis when the pH is reduced, they hydrolyze to release their component fragrance compounds when applied to skin or are exposed even to reduced pH such as present in air and humidity. The reduction in pH should be at least 0.1, preferably at least about 0.5 units. Preferably the pH is reduced by at least 0.5 units to a pH of 7.5 or less, more
20 preferably 6.9 or less. Preferably, the solution in which the pro-accord is applied is alkaline.

Odor Value

The pro-fragrances of the present invention typically have an Odor Value greater than or equal to about 1, preferably greater than or equal to about 5, more preferably greater than or equal to about 10. The term "Odor Value" is defined by the following formula:
25

$$\text{OV} = \frac{[\text{Concentration of FRM}]}{\text{ODT}}$$

wherein OV is the odor value of the fragrance raw material released upon the skin by the pro-accord. The odor value is the concentration of the fragrance raw material, FRM, on the skin
30 surface divided by the Odor Detection Threshold, ODT. The term "level of noticeability" is often applied to and/or substituted for the term "odor value".

Odor Detection Threshold

For the purposes of the present invention the term "odor detection threshold" is defined as the level at which a fragrance raw material is perceptible to the average human. The odor

detection threshold (ODT) of the compositions of the present invention are preferably measured by carefully controlled gas chromatograph (GC) conditions as described hereinbelow. The preferred fragrance raw materials of the present invention have an ODT of at least about 100 part per billion (ppb), more preferably 10 ppb, most preferably 1 ppb. Fragrance raw materials
5 having an ODT greater than 10 parts per million (ppm) are typically avoided unless useful as an adjunct ingredient, for example, as an adjunct alcohol when adjusting the fragrance release half-life of an orthoester.

Determination of Odor Detection Thresholds is as follows. A gas chromatograph is characterized to determine the exact volume of material injected by a syringe, the precise split
10 ratio, and the hydrocarbon response using a hydrocarbon standard of known concentration and chain-length distribution. The air flow rate is accurately measured and, assuming the duration of a human inhalation to last 0.02 minutes, the sampled volume is calculated. Since the precise concentration at the detector at any point in time is known, the mass per volume inhaled is known and hence the concentration of material. To determine whether a material has a threshold below
15 10 ppb, solutions are delivered to the sniff port at the back-calculated concentration. A panelist sniffs the GC effluent and identifies the retention time when odor is noticed. The average over all panelists determines the threshold of noticeability or ODT. The necessary amount of analyte is injected onto the column to achieve a 10 ppb concentration at the detector. Typical gas chromatograph parameters for determining odor detection thresholds are listed below.

20 GC: 5890 Series II with FID detector 7673 Autosampler

Column: J&W Scientific DB-1, length 30 m, i.d. 0.25 mm, film thickness 1 μ m.

Split Injection: 17/1 split ratio

Autosampler: 1.13 μ l/injection

Column flow: 1.10 mL/min

25 Air flow: 345 mL/min

Inlet temperature: 245° C

Detector temperature: 285° C

Temperature Information:

Initial temperature: 50° C

30 Rate: 5° C/min

Final temperature: 280° C

Final time: 6 min

Leading assumptions: 0.02 minutes per sniff and that GC air adds to sample dilution.

Skin Performance Index

Although a pro-fragrance or pro-accord may comprise a fragrance release half-life which ensures delivery of a fragrance raw material during a period of time useful to the formulator, unless the fragrance raw materials which comprise said fragrance delivery system have ODT values large enough to be perceived by the user, the formulator will be compelled to use an inordinate amount of material to achieve a suitable fragrance level.

The pro-fragrances of the present invention have a Skin Performance Index (SPI) greater than or equal to 0.1, preferably greater than or equal to 0.5. The Skin Performance Index is defined by the following:

$$\text{SPI} = \frac{[\text{Odor Value}]^*}{t_{1/2}}$$

wherein the term [Odor Value]* is the estimated concentration of the fragrance raw material in the headspace above a solution of the fragrance raw material as measured in a 1% solution of ethanol, and $t_{1/2}$ is the fragrance release half-life measured at pH 5.3 in the above described buffer. For the purposes of the present invention, the $t_{1/2}$ of the SPI is measured at 5.3 and the value of the fragrance release half-life is preferably from 0.1 hours to 60 hours.

The [Odor Value]* is an estimation of the vapor pressure of the fragrance raw material using empirically determined KOVATS indices. "The Vapor Pressures of Pure Substances", T. Boublik et al., Elsevier, New York (1973) incorporated herein by reference, describes an index line for normal alkanes wherein C_{10} is equal to 30,000 ppb, C_{12} is equal to 3,000 ppb, C_{14} is equal to 300 ppb, C_{16} is equal to 30 ppb, etc. Using these values as reference standards, the KOVATS index of a fragrance raw material is obtained from gas chromatographic analysis of the FRM and the experimental index is then used to determine the relative vapor pressure and hence the head space concentration of the fragrance raw material.

"New Method for Estimating Vapor Pressure by the Use of Gas Chromatography" *J. Chromatography A*, 79 p 123-129, (1996) and "Simple and Versatile Injection System for Capillary Gas Chromatographic Columns: Performance Evaluation of a System Including Mass Spectrometric and Light Pipe Fourier-Transform Infrared Detection", *J. Chromatography A*, 713, p 201-215, (1996) included herein by reference, further describe methods and techniques suitable for use in determining the vapor pressure and head space concentration of FRM's as they relate to the term [Odor Value]* of the present invention.

Using the criteria set forth in the present invention *inter alia* fragrance release half-life, odor value, odor detection threshold, skin performance index, the formulator is able to fashion an aldehyde or ketone releasing cyclic pro-fragrance. By manipulation of the R^2 , R^3 , R^4 , R^5 and R^6

units of the cyclic pro-fragrances of the present invention, the release rate of either an aldehyde or ketone fragrance raw material can be adjusted. Several different pro-fragrances which release the same fragrance raw material, but at differing rates or levels, can be admixed to further prolong or extend the period of fragrance raw material delivery.

5 When present, the carriers, fixatives, or stabilizers will comprise the balance of the compositions. Typical carriers are methanol, ethanol (preferred), iso-propanol, polyethylene glycol, as well as water in some instances, especially as a vehicle to deliver materials which provide reserve alkalinity to the fragrance delivery system. Fixatives serve to lower the volatility of certain top and middle notes in order to extend their contact time on skin. Adjunct ingredients
10 include perfume raw material components which are essential oils and are therefore not a single chemical entity. In addition, the adjunct ingredients may be mixtures of materials which serve a purpose in addition to providing a pleasurable odor (i.e., an astringent in a personal hygiene article).

 The formulators, in addition to selected fragrance raw materials, can combine a pre-
15 blended fragrance with the pro-fragrances of the present invention. For the purposes of the present invention the term "pre-blended fragrance" is defined as an existing fragrance accord, commercially available or otherwise, which is enhanced by or in turn enhances the fragrance delivered by the pro-fragrances and/or pro-accords which comprise the balance of the fragrance delivery system. For example, an existing fragrance accord may be suitable in the short term but
20 due to differential evaporation of the most volatile components, high notes, *inter alia*, the fragrance diminishes or the bouquet shifts. Therefore, pro-fragrances and pro-accords are combined which will release said volatile components thereby sustaining the complete fragrance accord.

 The following is a non-limiting example of a β -amino ketone pro-fragrance according to
25 the present invention.

EXAMPLE 1

1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N-(2-hydroxyethyl)-N-phenylmethyl-1-butanone

 The damascone releasing β -amino ketone pro-fragrance adduct is prepared by stirring 1 equivalent of δ damascone with 2 equivalents of *N*-benzylethanolamine until complete as
30 indicated suitable analytical method *inter alia*, thin layer chromatography, ^1H NMR. The product is purified by column chromatography on Aluminum Oxide (activated, basic, Brockman I, standard grade, 150 mesh, 58 angstrom) using a mobile phase consisting of 50% hexane, 49% chloroform and 1% triethylamine. Solvents are removed *in vacuo* and the product is placed

under high vacuum (0.10 mm Hg) for 24 hours with stirring to remove any residual solvent to yield the desired β -amino ketone pro-fragrance.

The following are examples of fine fragrance compositions which comprise the fragrance delivery system of the present invention.

TABLE I

Ingredients	Weight %			
	2	3	4	5
β -Amino ketone pro-fragrance ¹	0.4	0.8	--	--
β -Amino ketone pro-fragrance ²	--	--	0.3	0.6
Pro-fragrance ³	0.2	0.4	--	--
Pro-fragrance ⁴	0.2	0.4	--	--
Pro-fragrance ⁵	--	--	0.6	0.2
Pro-fragrance ⁶	--	--	--	0.2
Pre-formulated accords ⁷	14.0	14.0	14.0	14.0
Fragrance raw material ⁸	1.98	1.98	--	--
Fragrance raw material ⁹	0.02	0.02	--	--
Carrier ¹⁰	balance	balance	balance	balance


1. 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone.
2. 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(2-hydroxyethyl)-1-butanone.
3. 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester.
4. 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester.
5. Tris(geranyl) orthoformate.
6. Benzyl bis(geranyl) orthoformate.
7. Admixture of one or more fragrance raw materials having a pro-fragrance enhancing of compatible bouquet.
8. Lillial.
9. γ -decalactone.
10. Ethanol.

What is claimed is:

1. A fragrance raw material delivery system comprising:


A) from 0.01% by weight, of a pro-fragrance component comprising:

a) at least 0.01% by weight, of an β -amino ketone pro-fragrance component, said pro-fragrance having the formula:




wherein G^1 and G^2 are methyl, or a unit selected from the group consisting of:


i) 2,6,6-trimethylcyclohex-2-enyl having the formula:




ii) 2,6,6-trimethylcyclohex-1-enyl having the formula:



iii) 2-methylene-6,6-dimethylcyclohexanyl having the formula:



iv) 2,6,6-trimethylcyclohex-3-enyl having the formula:




v) and mixtures thereof;

provided at least one G^1 or G^2 is methyl and further provided both G^1 and G^2 are not both methyl; R and R^1 are each independently C_1 - C_{22} substituted or unsubstituted, branched or unbranched alkyl, C_2 - C_{22} substituted or unsubstituted, branched or unbranched alkenyl, C_2 - C_{20} substituted or unsubstituted, branched or unbranched hydroxyalkyl, C_7 - C_{20} substituted or unsubstituted alkylenearyl, C_3 - C_{20} substituted or unsubstituted cycloalkyl, alkyleneoxy units having the formula:



wherein R^4 is C_2 - C_4 alkylene, R^5 is hydrogen, C_1 - C_4 alkyl, and mixtures thereof, x is from 1 to 6; C_6 - C_{20} aryl, C_5 - C_{20} heteroaryl comprising one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof; two of the units Y^1 , Y^2 , R or R^1 can be taken together to form one or more aromatic or non-aromatic, heterocyclic or non-heterocyclic, single rings, fused rings, bicyclo rings, spiroannulated rings, or mixtures thereof, said rings comprising from 3 to 20 carbon atoms and one or more heteroatoms selected from the group consisting of nitrogen, oxygen, sulfur, and mixtures thereof;

b) at least 0.01% by weight, of an aldehyde or ketone releasing pro-fragrance component, said pro-fragrance having the formula:



wherein said pro-fragrance or pro-acord releases an aldehyde or a ketone fragrance raw material, wherein X is oxygen or sulfur; R is:

- i) C_6 - C_{22} substituted or unsubstituted linear alkyl;
- ii) C_6 - C_{22} substituted or unsubstituted branched alkyl;
- iii) C_6 - C_{22} substituted or unsubstituted linear alkenyl;

- iv) C₆-C₂₂ substituted or unsubstituted branched alkenyl;
 v) C₆-C₂₂ substituted or unsubstituted cycloalkyl;
 vi) C₆-C₂₂ substituted or unsubstituted branched cycloalkyl;
 vii) C₆-C₂₂ substituted or unsubstituted cycloalkenyl;
 5 viii) C₆-C₂₂ substituted or unsubstituted branched cycloalkenyl;
 ix) C₆-C₂₂ substituted or unsubstituted aryl;
 x) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
 xii) and mixtures thereof;
- 10 R¹ is:
 i) hydrogen;
 ii) C₁-C₁₀ substituted or unsubstituted linear alkyl;
 iii) C₃-C₁₀ substituted or unsubstituted branched alkyl;
 iv) C₂-C₁₀ substituted or unsubstituted linear alkenyl;
 15 v) C₃-C₁₀ substituted or unsubstituted branched alkenyl;
 vi) C₃-C₁₅ substituted or unsubstituted cycloalkyl;
 vii) C₄-C₁₅ substituted or unsubstituted branched cycloalkyl;
 viii) C₄-C₁₅ substituted or unsubstituted cycloalkenyl;
 ix) C₅-C₁₅ substituted or unsubstituted branched cycloalkenyl;
 20 x) C₆-C₁₅ substituted or unsubstituted aryl;
 xi) C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 xii) C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
- R and R¹ can be taken together to form a substituted or unsubstituted ring
 having in the ring from 3 to 10 carbon atoms; and
 25 each R², R³, R⁶ and each R⁴ and R⁵ pair are independently:
 i) R¹;
 ii) hydroxyl;
 iii) a carbonyl comprising unit having the formula:
- $$\text{---}(\text{CH}_2)_x\text{COR}^7$$
- 30 wherein R⁷ is:
 a) -OH;
 b) -OR⁸ wherein R⁸ is hydrogen, C₁-C₁₅ substituted or
 unsubstituted linear alkyl, C₁-C₁₅ substituted or
 unsubstituted branched alkyl, C₂-C₂₂ substituted or

- 5
- 10
- 15
- iv) unsubstituted linear alkenyl, C₃-C₂₂ substituted or unsubstituted branched alkenyl, or mixtures thereof;
 -N(R⁹)₂ wherein R⁹ is hydrogen, C₁-C₆ substituted or unsubstituted linear alkyl, C₃-C₆ substituted or unsubstituted branched alkyl, or mixtures thereof;
 C₁-C₂₂ substituted or unsubstituted linear alkyl;
 C₁-C₂₂ substituted or unsubstituted branched alkyl;
 C₂-C₂₂ substituted or unsubstituted linear alkenyl;
 C₃-C₂₂ substituted or unsubstituted branched alkenyl;
 C₃-C₂₂ substituted or unsubstituted cycloalkyl;
 C₆-C₂₂ substituted or unsubstituted aryl;
 C₆-C₂₂ substituted or unsubstituted heterocyclicalkyl;
 C₆-C₂₂ substituted or unsubstituted heterocyclicalkenyl;
 the index x is from 0 to 22;
- iv) alkyleneoxy units having the formula:



wherein each R¹⁰, R¹¹, and R¹² is independently;

- 20
- a) hydrogen;
 b) -OH;
 c) C₁-C₄ alkyl;
 d) or mixtures thereof;

R¹³ is:

- 25
- a) hydrogen;
 b) C₁-C₄ alkyl;
 c) or mixtures thereof;

R¹⁴ is:

- 30
- a) hydrogen;
 b) C₁-C₄ alkyl;
 c) or mixtures thereof;

R¹⁰ and R¹¹ can be taken together to form a C₃-C₆ spiroannulated ring, carbonyl unit, or mixtures thereof; y has the value from 0 to 10, z has the value from 1 to 50;

- v) any two R², R³, R⁴, R⁵, or R⁶ units can be taken together to form:


- 5
- a) a carbonyl moiety;
 b) a C₃-C₆ spiroannulated ring;
 c) a heterocyclic aromatic ring comprising from 5 to 7 atoms;
 d) a non-heterocyclic aromatic ring comprising from 5 to 7 atoms;
 e) a heterocyclic ring comprising from 5 to 7 atoms;
 f) a non-heterocyclic ring comprising from 5 to 7 atoms;
 g) or mixtures thereof;
- 10 vi) and mixtures thereof; and
 the index n is an integer from 1 to 3; and
- B) optionally from 1% by weight, a fragrance raw material component comprising:
- i) optionally at least 1% by weight, of a mixture of one or more base note fragrances;
- 15 ii) optionally at least 1% by weight, of a mixture of one or more top or middle note fragrances;
- iii) optionally the balance carriers, fixatives, and other adjunct ingredients.
2. A composition according to Claim 1 wherein R is C₁-C₈ alkyl, benzyl, and mixtures thereof; R¹ is hydrogen, C₁-C₈ alkyl, C₁-C₄ hydroxyalkyl, alkyleneoxy units having the formula:
- 20
$$-(R^4O)_xR^5$$
- wherein R⁴ is C₂-C₄ alkylene, R⁵ is hydrogen, C₁-C₄ alkyl, and mixtures thereof, x is from 1 to 6; and mixtures thereof.
- 25
3. A composition according to either Claim 1 or 2 wherein said aldehyde or ketone releasing pro-fragrance component releases a fragrance raw material selected from the group consisting of 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde, phenylacetaldehyde, methylnonyl acetaldehyde, 2-phenylpropan-1-al, 3-phenylprop-2-en-1-al, 3-phenyl-2-pentylprop-2-en-1-al, 3-phenyl-2-hexylprop-2-enal, 3-(4-isopropylphenyl)-2-methylpropan-1-al, 3-(4-ethylphenyl)-2,2-dimethylpropan-1-al, 3-(4-tert-butylphenyl)-2-methyl-propanal, 3-(3,4-methylenedioxyphenyl)-2-methylpropan-1-al, 3-(4-ethylphenyl)-2,2-dimethylpropanal, 3-(3-isopropylphenyl)butan-1-al, 2,6-dimethylhept-5-en-1-al, n-decanal, n-undecanal, n-dodecanal, 3,7-dimethyl-2,6-octadien-1-al, 4-methoxybenzaldehyde, 3-methoxy-4-hydroxybenz-aldehyde, 3-ethoxy-4-
- 30
- 35

hydroxybenzaldehyde, 3,4-methylenedioxy-benzaldehyde, 3,4-dimethoxybenzaldehyde, and mixtures thereof.

4. A composition according to any of Claims 1-3 further comprising at least 0.01% by weight, of a pro-accord which releases $n + 1$ fragrances raw materials wherein n is the number of fragrance raw materials from which said pro-accord is formed, n is from 1 to 3.
5. A fragrance delivery system comprising:
 - a) from 0.1% by weight, of a β -amino ketone pro-fragrance selected from the group consisting of 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(2-hydroxyethyl)-1-butanone, and mixtures thereof;
 - b) from 0.2% by weight, of an aldehyde releasing pro-fragrances selected from the group consisting of 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, and mixtures thereof;
 - c) optionally from 0.1% by weight, of one or more alcohol releasing pro-fragrances or pro-accords;
 - d) from 0.1% by weight, of one or more fragrance raw materials;
 - e) optionally, from 1% by weight, of pre-blended perfume ingredients of fragrance raw material accords; and
 - f) the balance carriers.
6. A system according to Claim 5 wherein said alcohol releasing pro-fragrances or pro-accords are selected from the group consisting of tris-geranyl orthoformate, tris(*cis*-3-hexen-1-yl) orthoformate, tris(phenylethyl) orthoformate, bis(citronellyl) ethyl orthoacetate, tris(citronellyl) orthoformate, tris(*cis*-6-nonenyl) orthoformate, tris(phenoxyethyl) orthoformate, tris(geranyl, neryl) orthoformate (70:30 geranyl:neryl), tris(9-decenyl) orthoformate, tris(3-methyl-5-phenylpentanyl) orthoformate, tris(6-methylheptan-2-yl) orthoformate, tris([4-(2,2,6-trimethyl-2-cyclohexen-1-yl)-3-buten-2-yl] orthoformate, tris[3-methyl-5-(2,2,3-trimethyl-3-cyclopenten-1-yl)-4-penten-2-yl] orthoformate, trismenthyl orthoformate, tris(4-isopropylcyclohexylethyl-2-yl) orthoformate, tris-(6,8-dimethylnonan-2-yl) orthoformate, tris-phenylethyl orthoacetate, tris(*cis*-3-hexen-1-yl) orthoacetate, tris(*cis*-6-nonenyl) orthoacetate, tris-citronellyl

orthoacetate, bis(geranyl) benzyl orthoacetate, tris(geranyl) orthoacetate, tris(4-isopropylcyclohexylmethyl) orthoacetate, tris(benzyl) orthoacetate, tris(2,6-dimethyl-5-heptenyl) orthoacetate, bis(*cis*-3-hexen-1-yl) amyl orthoacetate, and neryl citronellyl ethyl orthobutyrate, and mixtures thereof.


- 5
7. A system according to either of Claims 5 or 6 wherein said alcohol releasing pro-fragrances or pro-accords are selected from the group consisting of bis(ethyl) bis(geranyl) orthocarbonate, bis(ethyl) bis(phenylethyl) orthocarbonate, bis(ethyl) bis(*cis*-3-hexenyl) orthocarbonate, bis(ethyl) bis(citronellyl) orthocarbonate, bis(ethyl) bis(linalyl) orthocarbonate, bis(ethyl) bis(menthyl) orthocarbonate, bis(dodecyl) bis(geranyl) orthocarbonate, and bis(dodecyl) bis(phenylethyl) orthocarbonate, and mixtures thereof.
- 10
8. A fine fragrance or perfume comprising:
- 15
- a) from 0.1% by weight, of a β -amino ketone pro-fragrance selected from the group consisting of 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(5-hydroxy-3-oxapentyl)-1-butanone, 1-(2,6,6-trimethyl-3-cyclohexen-1-yl)-3-N,N-bis(2-hydroxyethyl)-1-butanone, and mixtures thereof;
- 20
- b) from 0.2% by weight, of an aldehyde releasing pro-fragrances selected from the group consisting of 2-(6-methyl-5-hepten-2-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, 2-(2,4-dimethyl-3-cyclohexen-1-yl)-3-(1-methylethyl)-4-oxazolidinecarboxylic acid methyl ester, and mixtures thereof;
- c) optionally from 0.1% by weight, of one or more orthoester pro-accords; and
- d) the balance carriers.
- 25
9. A composition according to Claim 8 wherein said pro-accords have the formula:



wherein R is hydrogen, C₁-C₈ linear alkyl, C₄-C₂₀ branched alkyl, C₆-C₂₀ cyclic alkyl, C₆-C₂₀ branched cyclic alkyl, C₆-C₂₀ linear alkenyl, C₆-C₂₀ branched alkenyl, C₆-C₂₀ cyclic alkenyl, C₆-C₂₀ branched cyclic alkenyl, C₆-C₂₀ substituted or unsubstituted aryl, and mixtures thereof; R¹, R² and R³ are independently C₁-C₂₀ linear, branched, or substituted alkyl; C₂-C₂₀ linear, branched, or substituted alkenyl; C₅-C₂₀ substituted or

30

5 unsubstituted cyclic alkyl; C₆-C₂₀ substituted or unsubstituted aryl, C₂-C₄₀ substituted or unsubstituted alkyleneoxy; C₃-C₄₀ substituted or unsubstituted alkyleneoxyalkyl; C₆-C₄₀ substituted or unsubstituted alkylenearyl; C₆-C₃₂ substituted or unsubstituted aryloxy; C₆-C₄₀ substituted or unsubstituted alkyleneoxyaryl; C₆-C₄₀ oxyalkylenearyl, and mixtures thereof; or a cyclic orthoester having the formula:



10 wherein at least one R⁴ or R⁵ is derived from a fragrance raw material alcohol, each X is -C(R⁶)₂- wherein each R⁶ is independently hydrogen, C₁-C₂₂ linear or branched alkyl, C₂-C₂₂ linear or branched alkenyl, C₆-C₂₂ substituted or unsubstituted aryl, and mixtures thereof, Y is -CR⁷R⁸-, C=O, and mixtures thereof, wherein R⁷ and R⁸ are independently hydrogen, hydroxyl, nitro, nitrilo, C₁-C₃₀ substituted or unsubstituted linear alkyl, C₃-C₃₀ substituted or unsubstituted branched alkyl, C₃-C₃₀ substituted or unsubstituted cyclic alkyl, C₂-C₃₀ substituted or unsubstituted linear alkenyl, C₃-C₃₀ substituted or unsubstituted branched alkenyl, C₃-C₃₀ substituted or unsubstituted cyclic alkenyl, C₂-C₃₀ substituted or unsubstituted linear alkynyl, C₃-C₃₀ substituted or unsubstituted branched alkynyl, C₆-C₃₀ substituted or unsubstituted alkylenearyl, C₆-C₃₀ substituted or unsubstituted aryl, C₂-C₂₀ substituted or unsubstituted alkyleneoxy, C₃-C₂₀ substituted or unsubstituted alkyleneoxyalkyl, C₇-C₂₀ substituted or unsubstituted alkylenearyl, C₆-C₂₀ substituted or unsubstituted alkyleneoxyaryl, and mixtures thereof, or R⁷ and R⁸ can be taken together to form a spiroannulated ring or taken together with any R⁶ to form a fused ring, said spiroannulated or fused ring having from 3 to 8 carbons and optionally one or more heteroatoms in said ring, said ring further optionally substituted by one or more C₁-C₂₂ alkyl, C₁-C₂₂ alkenyl, C₆-C₁₂ aryl, C₆-C₂₂ alkylenearyl units, and mixtures thereof; m is from 0 to 14, p is from 0 to 14, and n is from 0 to 3; provided m + n + p is at least 1 and less than or equal to 14.

10. A composition according to either of Claims 8 or 9 wherein said orthoester pro-accord is a pro-accord which comprises *n* fragrance raw materials, said fragrance raw materials having a molecular weight greater than or equal to 100 g/mol and capable of releasing upon hydrolysis *n* + 1 fragrance raw materials, provided said pro-accord:
- 30 a) has a molecular weight greater than or equal to 300 g/mol;

- b) has a molecular weight at least two times greater than the lowest molecular weight fragrance raw material which comprises said pro-accord; and
 - c) has a fragrance release half-life of greater than or equal to 0.1 hours at pH 5.3 and less than or equal to 12 hours at pH 2.5 when measured in NaH_2PO_4 buffer;
- 5 wherein n is an integer from 1 to 3.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/10166

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D3/50 A61K7/46 C07C225/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D A61K C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 98 47477 A (PROCTER & GAMBLE) 29 October 1998 (1998-10-29) page 22 -page 29; claims	1,5-10
A	WO 99 00347 A (SIVIK MARK ROBERT ;PROCTER & GAMBLE (US)) 7 January 1999 (1999-01-07) claims 1-5,9	1,5
A	US 4 129 569 A (SCHREIBER WILLIAM L ET AL) 12 December 1978 (1978-12-12) the whole document	1,5
A	US 4 011 233 A (DUBS PAUL ET AL) 8 March 1977 (1977-03-08) the whole document	1,5

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

28 July 2000

Date of mailing of the international search report

18.08.00

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Grittern, A

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 00/10166

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9847477 A	29-10-1998	US 6077821 A AU 7139298 A EP 0977548 A	20-06-2000 13-11-1998 09-02-2000
WO 9900347 A	07-01-1999	EP 0991615 A	12-04-2000
US 4129569 A	12-12-1978	GB 1601729 A JP 54005970 A	04-11-1981 17-01-1979
US 4011233 A	08-03-1977	NONE	

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 00/10166

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.: 1-10 (partial)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-10 (partial)

The present application claims pro-fragrances, comprising i.a. beta-amino ketones having the formula as it is disclosed in claim 1 of the application. The meaning of substituents R2 and R3 is not explained.

Therefor the search has been restricted to the examples as been disclosed on page 9, lines 1-10 of the description.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.